

Exponential Bounds for Random Sums.

MIGDASHIEV B.M.^b, OSTROVSKY E.I.^{a* 1 2}

Abstract. We construct a non - improved exponential bounds for distribution of normed sums of i.i.d. random variables with random numbers of summand.

Key words: Random sum, exponential estimation, Orlicz spaces, martingales, saddle - point method.

1. Introduction.

Let (Ω, F, μ) be a probability space, $\{\xi(i)\}, i = 1, 2, \dots$ be a sequence of independent identical distributed (i.i.d) centered: $\mathbf{E}\xi(i) = 0$ random variables (r.v) with finite non - trivial variance $\sigma^2 = \mathbf{D}\xi(i) \in (0, \infty)$, and let $\eta, \eta \geq 1$ be an integer r.v. with finite first moment $\mathbf{E}\eta = A$, where $A \in [2, \infty)$. We assume at first that the r.v. η and the sequence $\{\xi(i)\}$ are independent.

We will denote for arbitrary r.v. τ and $x = \text{const} \geq 0$ the tail function

$$T(\tau, x) = \max(\mathbf{P}(\tau \geq x), \mathbf{P}(\tau \leq -x)),$$

will write for the r.v. $\xi(1)$ $R(x) = T(\xi(1), x)$, and we define the so - called normed random sum and corresponding uniform tail function

$$S = \sum_{i=1}^{\eta} \xi(i) / (\sigma \sqrt{A}),$$

$$V(x) = V(\text{Law}(\eta), \text{Law}(\xi(i)), x) = \sup_{A \geq 2} T(S, x).$$

In the case if the r.v. $\eta - 1$ has a Poisson distribution $\text{Pois}(A)$ with parameter A : $\mathbf{P}(\eta - 1 = n) = A^n \exp(-A)/n!$, $n = 0, 1, 2, \dots$ and $\text{Law}(\eta - 1) \in \cup_{A \geq 2} \{\text{Pois}(A)\}$ we will write $V(\text{Pois}; \text{Law}(\xi(i)), x) = V(\cup_{A \geq 2} \{\text{Pois}(A)\}, \text{Law}(\xi(i)), x)$; if the r.v. η has a geometrical distribution $G(A)$: $\mathbf{P}(\eta = n) = A^{-1}(1 - 1/A)^{n-1}$, $n = 1, 2, \dots$, $A \geq 2$ and $\text{Law}(\eta) \in \cup_{A \geq 2} \{G(A)\}$ we will write $V(G; \text{Law}(\xi(i)), x) =$

¹Corresponding Author.

²Department of Mathematics and Computer Science, Ben - Gurion University, Beer - Sheva, 84105, Ben - Gurion street, 4; P.O. Box 61, Israel.
E - mail: galaostr@cs.bgu.ac.il

$V(\cup_{A \geq 2}\{G(A)\}, \text{Law}(\xi(i)), x)$; for the case of *all* distribution r.v. η under the condition that $\mathbf{E}\eta = A$, $\exists A \geq 2$ we will use the notations correspondently $\text{Dis}(A)$, $\text{Dis} = \cup_{A \geq 2}\{\text{Dis}(A)\}$ and $V(\text{Dis}, \text{Law}(\xi(i)), x) = V(\cup_{A \geq 2}(\{\text{Dis}(A)\}), \text{Law}(\xi(i)), x)$.

Our goal is the bidirectional exponential estimating $V(x)$ at $x \rightarrow \infty$, $x \geq 2$ in the terms of distributions $\text{Law}(\xi(i))$, $\text{Law}(\eta)$.

We have for all distribution $\xi(i)$ under the conditions $\mathbf{E}\xi(i) = 0$, $\mathbf{D}\xi(i) \in (0, \infty)$, since $\mathbf{E}\xi(i) = 0$, $\mathbf{D}\xi(i) \in (0, \infty)$:

$$V(\text{Dis}, \text{Law}(\xi(i)), x) \leq \min(1, x^{-2})$$

by virtue of Chebyshev inequality; this estimation is called trivial.

There are many publications about the moment estimations $\mathbf{E}|S|^p$ and statistical applications of those estimations (see, for example, (Gut A., 1988), (Gut A., 2003):

$$|S|_p \leq B(p) |\eta|_p^{1/2} |\xi(1)|_p, \quad p \geq 2, \quad (0)$$

where $B(p)$ is a constant in the famous Burkholder inequality for martingales. Here and further for arbitrary r.v. ζ

$$|\zeta|_p = \mathbf{E}^{1/p}|\zeta|^p = \|\zeta\|_{L_p(\Omega, \mathbf{P})}.$$

It is proved in (Hitczenko, 1990) that the best boundary for $B(p)$ at $p \geq 2$, $p \rightarrow \infty$ is $B(p) \leq C p / \log p$, C is an *absolute* constant. Note that the estimation (0) is proved in the case if r.v. η is the *stopping time* for the sequence $\{\xi(i)\}$, for example, if $\forall n = 1, 2, \dots$ r.v.'s $\xi(1), \xi(2), \dots, \xi(n)$ and the event $\{\eta = n\}$ are independent.

See also (Gine at al., 2003) etc. For non - random sums, i.e. in the case $\mathbf{P}(\eta = n) = 1$ for some $n = 1, 2, \dots$ the exponential bounds for $V(x)$ are constructed in (Buldygin at al., 1992), (Ostrovsky, 1999, p. 28); for more generally case if $\{\xi(i)\}$ are the martingale - differences the exponential bound for $T(S, x)$ is derived in (Lesign at al., 2001).

For reader convenience we recall here some results of (Buldygin at al., 1992), (Lesign at al., 2001). Let us define for some tail function $T(x)$, i.e. for the function $T = T(x)$ under the conditions: $T(0) = 1$, $T(\infty) = 0$, monotonically non - increasing and right continuous with finite second moment $|\int_0^\infty x^2 dT(x)| < \infty$ the operator

$$W[T](x) = \min\left(1, 4 \inf_{z > 0} \left[\exp(-x^2/(8z^2)) - \int_z^\infty x^2 dT(x)\right]\right).$$

Further, put

$$\varphi(\lambda) = \max_{\pm} \log \mathbf{E} \exp(\pm \lambda \xi(i)).$$

This definition is non - trivial only if the variable $\xi(1)$ satisfies the so - called Kramer condition:

$$\exists \lambda_0 \in (0, \infty), \forall \lambda \in (-\lambda_0, \lambda_0) \Rightarrow \varphi(\lambda) < \infty;$$

in other case we set $\varphi(\lambda) = \infty \forall \lambda \neq 0$.

Let us introduce the function $\chi(\lambda) = \sup_{n=1,2,\dots} n\varphi(\lambda/\sqrt{n})$,

$$\chi^*(x) = \sup_{\lambda} (\lambda x - \chi(\lambda)),$$

$$Q[R](x) = \min\{W[R](x), \exp(-\chi^*(x))\}.$$

Lemma 1. (Buldygin at al., 1992), (Lesign at al., 2001).

$$\sup_n T \left(n^{-1/2} \sum_{i=1}^n \xi(i), x \right) \leq Q[R](x), \quad x \geq 0. \quad (1)$$

Let us introduce the Orlicz spaces $G(m, r)$ (in order to describe the examples) of random variables as the set of all r.v. $\{\tau\}$ with finite norm

$$\|\tau\|_{m,r} = \sup_{p \geq 2} |\tau|_p p^{-1/m} \log^{r/m} p.$$

Here $m = \text{const} > 0$, $r = \text{const} \in R^1$. It is easy to verify that $G(m, r)$ is isomorphic to the Orlicz space with N - function

$$N(u) = \exp(|u|^m \log^r |u|), \quad |u| \geq 2,$$

and that $\tau \in G(m, r)$ if and only if

$$T(\tau, x) \leq \exp(-C_1(m, r)x^m \log^r(C_2(m, r) + x)). \quad (2)$$

See (Buldygin at al, 1992, p.351).

For example, assume that the r.v. τ has Poisson distribution with parameter A ; $A \geq 2$. Then for some non - trivial positive absolute constants C_1, C_2 and all $p, x \geq 2$

$$|\tau - A|_p \leq C_1 \sqrt{A} p / \log p,$$

or

$$\mathbf{P}(|\tau - A|/\sqrt{A} > x) \leq \exp(-C_2 x \log x).$$

Let us suppose, for example, that for some $m = \text{const} > 0, r \in R^1 \xi(1) \in G(m, r)$. We define the following functions $M = M(m, r), L = L(m, r)$: at $m \in (0, 1)$ or $m = 1, r < 0 \Rightarrow M = 2m/(m+2), L = 2r/(m+2)$; at $m = 1, r \geq 0$ or $m \in (1, 2), r < 0 \Rightarrow M = m, L = r$; at $m = 2, r \geq 0$ or $m > 2 \Rightarrow M = 2, L = 0$. We can define formally in the case $m = +\infty, r \in R \Rightarrow M = 2, L = 0$.

It follows from (1)

$$\sup_n T \left(\sum_{i=1}^n \xi(i)/\sqrt{n}, x \right) \leq \exp \left(-C_3 x^M \log^L(C_2 + x) \right), \quad (3)$$

$C_{2,3} = C_{2,3}(m, r)$, or, equally,

$$\sup_{n \geq 1} \left\| \sum_{i=1}^n \xi(i)/\sqrt{n} \right\|_{M,L} \leq C(m, r) \|\xi(1)\|_{m,r}.$$

It is proved in (Ostrovsky, 1999, p. 34) that in the case $m > 1$ the estimation (2) is exact at $x \rightarrow \infty$.

In this paper, the letter $C, C_j(\cdot)$ will denote positive finite various non-essentially constant which may differ from one formula to the next and which does not depend upon x, n . We make no attempt to obtain the best values for these constants.

2. Main result. Upper bound. Examples.

Theorem 1.

$$V(\text{Law}(\eta), \text{Law}(\xi(i)), \eta, x) \leq \mathbf{E} Q[R](\sigma x \sqrt{A/\eta}). \quad (4)$$

Proof. We will assume without loss of generality $\sigma = 1$. We receive from (1), using the formula of full probability and denoting $q_n = q_n(A) = \mathbf{P}(\eta = n)$: $V(x) =$

$$\begin{aligned} \mathbf{P} \left(\sum_{i=1}^n \xi(i) > x \sqrt{A} \right) &= \sum_{n=1}^{\infty} q_n \mathbf{P} \left(\sum_{i=1}^n \xi(i)/\sqrt{n} > x \sqrt{A/n} \right) \leq \\ &\sum_n q_n Q[R](x \sqrt{A/n}) = \mathbf{E} Q[R](x \sqrt{A/\eta}). \end{aligned}$$

Example 1. Let us suppose here that the r.v. η has a geometric distribution $G(A)$ with parameter $A, A \geq 2$, and that $\exists m > 0, \exists r \in R L(\xi(1)) \in G(m, r)$. We assert that at $x \geq 2 \Rightarrow \sup_{A \geq 2} V(G(A), G(m, r), x) \leq$

$$C_3(m, r) \exp \left(-C(m, r) x^{2M/(M+2)} (\log x)^{2L/(M+2)} \right)$$

$$\stackrel{def}{=} J(M, L; x) = J(x).$$

Proof. We have using theorem 1:

$$V(G(A), G(m, r), x) \leq \sum_{n=1}^{\infty} (A-1)^{-1} \exp(n[\log(1-1/A)] -$$

$$Cx^M n^{-M/2} A^{M/2} n^{-M/2} (\log^L(C_2 + x\sqrt{A/n})) \leq (A-1)^{-1} \times$$

$$C_1 \sum_{n=2}^{\infty} \exp\left(-C(n/A + x^M (A/n)^{M/2} (\log^L(C_2 + x\sqrt{A/n})))\right) \stackrel{def}{=}$$

$$\sum_{n=2}^{\infty} a(n; A, x).$$

We denote

$$N_0 = N_0(A, x) = \operatorname{argmax}_{n \geq 2} a(n; A, x) = A n_0(x),$$

where at $x \rightarrow \infty \Rightarrow$

$$N_0/A \sim C_2(M, L) x^{2M/(M+2)} (\log x)^{2L/(M+2)}.$$

Since the function $n \rightarrow a(n; A, x)$ is monotonic in the interval $[1, N_0]$ and $[N_0, \infty)$, we can estimate

$$C_2 (A-1) V(G, G(m, r), x) \leq \int_1^{\infty} \exp\left(-C\left((y/A) + x^M (\sqrt{A/y})^M \log^L(C_2 + x\sqrt{A/y})\right)\right) dy =$$

$$C_3 A \int_{1/A}^{\infty} \exp\left(-C\left(z - x^M z^{-M/2} \log^L(C_2 + x/\sqrt{z})\right)\right) dz \leq$$

$$C_3 A \int_0^{\infty} \exp\left(-C\left(z - x^M z^{-M/2} \log^L(C_2 + x/\sqrt{z})\right)\right) dz.$$

Let us denote $\beta = 2M/(2+M)$,

$$R(x, v) = v + (2/M) v^{-M/2} \log^L(C_2 + x^{2/(M+2)} v^{-1/2}),$$

$$U(x, v) = v + (2/M)v^{-M/2} \log^L x, \quad x \geq 2, v > 0;$$

$$S(x, v) = x^\beta \left(v + (2/M)v^{-M/2} \log^L (C_2 + x^{2/(M+2)}v^{-1/2}) \right).$$

Let $\Delta = \Delta(M, L)$ be the arbitrary function on M, L so that $\Delta > 2|L|/(M+2)$. After the substitution $z = x^\beta v$ we receive:
 $\sup_{A \geq 2} V(G(A), G(m, r), C_1 x)/C_2 \leq$

$$C x^\beta \int_0^\infty \exp(-S(x, v)) dv \stackrel{def}{=} I(M, L; x) = I(x).$$

We have $I = x^\beta(I_1 + I_2 + I_3)$, where

$$I_1 = \int_0^{\log^{-\Delta} x}, \quad I_2 = \int_{\log^{-\Delta} x}^{\log^{\Delta} x}, \quad I_3 = \int_{\log^{\Delta} x}^\infty \exp(-S(x, v)) dv.$$

Since at $v \in (0, \log^{-\Delta} x) \cup (\log^{-\Delta} x, \log^{+\Delta} x) \Rightarrow$

$$S(x, v) = x^\beta R(x, v) \geq C_3 U(C_4 x, v) \geq C x^\beta \log^{L+\Delta M/2} x,$$

we conclude

$$I_1 \leq \log^{-\Delta} x \cdot \exp\left(-C x^\beta \log^{L+\Delta M/2} x\right) \leq C_5 J(M, L; C_6 x);$$

$$x^\beta I_2 \leq (\log^{\Delta} x) \exp\left(-C \min_{\{v \in \log^{-\Delta} x, \log^{\Delta} x\}} U(x, v)\right) \leq$$

$C_7 J(M, N; C_8 x)$. Further,

$$x^\beta I_3 \leq \int_{\log^{\Delta} x}^\infty \exp(-x^\beta v) dv \leq C_9 J(M, N; C_{10} x).$$

Following, $V(G, G(m, r), x) \leq C_{11}(m, r) \times$

$$\exp\left(-C_{12}(m, r) x^{2M/(M+2)} (\log x)^{2L/(M+2)}\right), \quad x \geq 2. \quad (5)$$

On the other hand

$$\sup_{A \geq 2} \|S\|_{2M/(M+2), 2L/(M+2)} \leq C_6 \|\xi(1)\|_{m, r}.$$

Example 2. Let us now suppose again that $L\{\xi(i)\} \in G(m, r)$ for some $m > 0, r \in R$ and assume that the r.v. $\eta - 1$ has a *Poisson* distribution with parameter $B = A - 1$; $A \geq 2$. It follows from theorem 1 and Stirling's formula that $V(Pois(A), G(m, r), x) \leq$

$$\begin{aligned}
& C \sup_{B \geq 1} \sum_{n=1}^{\infty} \exp(-B + n \log B - \log n! - \\
& Cx^M B^{M/2} n^{-M/2} (\log^L(C_2 + x\sqrt{B/n}))) \leq \\
& C \sup_{B \geq 1} \sum_{n=1}^{\infty} \exp(-B - n \log n + \\
& n - Cx^M B^{M/2} n^{-M/2} (\log^L(C_2 + x\sqrt{B/n}))) \leq \\
& C \sum_{n=1}^{\infty} \sup_{B \geq 1} \exp(-B - n \log n + n - \\
& Cx^M B^{M/2} n^{-M/2} (\log^L(C_2 + x\sqrt{B/n}))). \tag{6}
\end{aligned}$$

It is easy to verify that the maximum of arbitrary member of the right - side (6) over $B \geq 1$ for sufficiently greater values $x \geq x_0$, $x_0 = \text{const} \geq 2$ is achieved at $B = 1$. Therefore $V(Pois(A), G(m, r), x)/C \leq$

$$\leq \sum_{n=2}^{\infty} \exp(-1 - n \log n + n - Cx^M n^{-M/2} \log^L(C_2 + x/\sqrt{n})).$$

Since for $x \geq 2$

$$\sum_{n \geq x^2} \exp(-n \log n + n) \leq \exp(-Cx^2 \log x),$$

we have:

$$V(Pois(A), G(m, r), x)/C \leq \exp(-Cx^2 \log x) +$$

$$\sum_{n \in [1, x^2]} \exp(n - n \log n - Cx^M n^{-M/2} \log^L(C_2 + x/\sqrt{n})).$$

Let us denote $N_1 = N_1(x) =$

$$\operatorname{argmax}_{n \in [1, x^2]} (n - n \log n - Cx^M n^{-M/2} \log^L(C_2 + x/\sqrt{n}));$$

it is easy to calculate:

$$N_1(x) \asymp C(M, L)x^{2M/(M+2)} (\log x)^{(2L-2)/(M+2)}, \quad x \rightarrow \infty.$$

We obtain for values $x \geq 2$:

$$\begin{aligned} V(Pois, G(m, r), x)/C(m, r) &\leq \exp(-Cx^2 \log x) + \\ (x^2 + 1) \exp\left(N_1 - N_1 \log N_1 - Cx^M N_1^{-M/2} \log^L(C_2 + x/\sqrt{N_1})\right) &\leq \\ C_3 \exp\left(-C(m, r)x^{2M/(M+2)} (\log x)^{(2L+M)/(M+2)}\right). \end{aligned} \quad (7)$$

In the case $m = \infty$, i.e. if the variable $\xi(1)$ is bounded (*mod* $\mathbf{P}$), then $2M/(M+2) = 1$, $(2L+M)/(M+2) = 1/2$.

We can rewrite (7) in the considered case $Law(\eta - 1) = Pois, Law(\xi(i) \in G(m, r))$ in the terms of $G(m, r)$ spaces:

$$\sup_{A \geq 2} \|S\|_{2M/(M+2), (M+2L)/(M+2)} \leq C(m, r) \|\xi(1)\|_{m, r}.$$

In the case if $Law\{\xi(i)\} \in \cup_{m>1}\{G(m, r)\}$ the estimation (7) improves some result of (Gine at al, 2003). For instance, if $m \in (1, \infty)$, $r = 0$, then from (7) it follows the inequality: $p \geq 2 \Rightarrow$

$$\sup_{A \geq 2} |S|_p \leq C_1 p^{1/2+1/\min(m,2)} / \sqrt{\log p}, \quad p \geq 2,$$

but we receive from (Gine at all, 2003):

$$\sup_{A \geq 2} |S|_p \leq C_2 p^{1+1/m} / \log p,$$

and we obtain from (Gut, 1988), (Gut, 2003):

$$\sup_{A \geq 2} |S|_p \leq C_3 p^{1+1/m} \log^{-1/2} p.$$

3. Low bounds. We will prove further that our estimations are non-improved in general case, for example, even for normal distribution of values $\{\xi(i)\}$.

Theorem 2. *We assert that for all values $m > 1$ and sufficiently larges $x, x \geq x_0 = const \geq 2$:* $V(G, G(m, r), x) \geq$

$$C_6(m, r) \exp\left(-C_7(m, r)x^{2M/(M+2)} (\log x)^{2L/(M+2)}\right), \quad (8)$$

$$V(Pois, G(m, r), x) \geq C_8(m, r) \exp \left(-C_9(m, r) x^{2M/(M+2)} (\log x)^{(2L+M)/(M+2)} \right). \quad (9)$$

Proof is very simple. It is enough to prove the inequality (8); the proposition (9) is proved analogously. Let $\xi(i)$ be independent symmetrically distributed r.v. with distributions

$$\mathbf{P}(|\xi(i)| > x) = \exp(-x^m [\log(C(m, r) + x)]^r), \quad x \geq 0,$$

and let us introduce the even smooth convex function $\varphi_{m,r}(\lambda) = \log \mathbf{E} \exp(\lambda \xi(i))$, $\lambda \in (-\infty, \infty)$. It is proved in (Buldygin et al., 1992, p.341), (Ostrovsky, 1999, p.34) that

$$C_1 \varphi_{m,r}(\lambda) \leq \psi_{m,r}(\lambda) \leq C_2 \varphi_{m,r}(\lambda),$$

where $\psi_{m,r}(\lambda) = \lambda^2$, $\lambda \in [-2, 2]$;

$$\psi_{m,r}(\lambda) = C_3 |\lambda|^{m/(m-1)} [\log(C_4 + |\lambda|)]^{-r/(m-1)}$$

in the case $|\lambda| > 2$. Since the r.v. $\{\xi(i)\}$ are i.i.d., we have for the non-random sum:

$$\log \mathbf{E} \exp \left(\lambda n^{-1/2} \sum_{i=1}^n \xi(i) \right) = n \varphi_{m,r}(\lambda/\sqrt{n}) \asymp$$

$n \psi_{m,r}(\lambda/\sqrt{n})$, where the symbol $\asymp$ is understood uniformly on n ; $\lambda \in R$:

$$0 < C_1(m, r) \leq \inf_{n \geq 1} \inf_{\lambda \in R} \frac{n \varphi_{m,r}(\lambda/\sqrt{n})}{n \psi_{m,r}(\lambda/\sqrt{n})} \leq \sup_{n \geq 1} \sup_{\lambda \in R} \frac{n \varphi_{m,r}(\lambda/\sqrt{n})}{n \psi_{m,r}(\lambda/\sqrt{n})} \leq C_2(m, r) < \infty.$$

We conclude at $x \geq 2$, $A \geq 2$:

$$\mathbf{P}(S > x) = \sum_{n=1}^{\infty} A^{-1} (1 - 1/A)^{n-1} \mathbf{P} \left(\sum_{i=1}^n \xi(i)/\sqrt{n} > x \sqrt{A/n} \right).$$

We deduce, choosing again in this sum only the member with $n = N_0$ (recall that $N_0 = N_0(x)$) and using the main result of paper (Bagdasarov et al., 1995):

$$\mathbf{P} \left(n^{-1/2} \sum_{i=1}^n \xi(i) > u \right) \geq \exp \left(-C_1 u^M \log^L(C_2 + u) \right),$$

where $C_1, C_2 = C_1(m, r), C_2(m, r)$; $u = u(n) \geq 2$:

$$\mathbf{P}(S > x) \geq (A - 1)^{-1} (1 - 1/A)^{N_0} \mathbf{P} \left(\sum_{i=1}^{N_0} \xi(i) / \sqrt{N_0} > x \sqrt{A/N_0} \right) \geq$$

$$C_6(m) \exp \left(-C_7(m) x^{2M/(M+2)} [\log x]^{2L/(M+2)} \right).$$

Theorem 3. For all values $x \geq 3$

$$V(Dis, N(0, 1), x) \geq C x^{-2}. \quad (10)$$

Here $N(0, 1)$ denotes the normal distribution with parameters 0, 1; and C is an absolute constant.

Proof. We suppose now $L(\xi(i)) = N(0, 1)$, i.e.

$$\mathbf{P}(\xi(i) > x) = \Psi(x) = (2\pi)^{-1/2} \int_x^\infty \exp(-y^2/2) dy.$$

We define $\alpha = 1/Ent[x^2]$, $Ent[z]$ denotes the integer part of z , for $x \geq 3$ and choose the r.v. η by the following way: $\mathbf{P}(\eta = 2) = 1 - 1/\alpha$, $\mathbf{P}(\eta = 1/\alpha) = \alpha$. Then $A = \mathbf{E}\eta = 3 - 2\alpha \geq 25/9 > 2$;

$$\mathbf{P}(S > x) > \alpha \mathbf{P} \left(\sum_{i=1}^{1/\alpha} \xi(i) > x \sqrt{A} \right) =$$

$$(1/[x^2]) \Psi \left(x \sqrt{3 - 2/[x^2]} \sqrt{1/[x^2]} \right) \geq x^{-2} \Psi \left(3\sqrt{3/8} \right) = Cx^{-2}.$$

4. Upper exponential bound for stopping time. In this section we will obtain the exponential bounds for the tails of distribution r.v. S in the case if η is the *stopping time* for the sequence $\{\xi(i)\}$, in addition to the moment estimations of S in (Gut, 1988), (Gut, 2003). Recall that again $\mathbf{E}S = 0$ and $\mathbf{D}S = 1$.

Theorem 4. Assume that the r.v. η belong to the space $G(m, r)$ for some $m > 0$, $r \in R$: $\eta \in G(m, r)$ and is the stopping time for the sequence $\{\xi(i)\}$, where $\xi(1) \in G(a, b)$, $a = \text{const} > 0, b = \text{const}$: $\forall x \geq 2$

$$T(\eta, x) \leq \exp(-C x^m \log^r x), \quad T(\eta, x) \leq \exp(-C x^a \log^b x).$$

Denote

$$q = \frac{2am}{2am + 2a + m}, \quad w = \frac{2arm + mb + 2am}{2am + 2a + m}.$$

We assert that at $x \geq 2$

$$\sup_{A \geq 2} \sup_{\eta: \mathbf{E}\eta = A} T(S, x) \leq \exp(-C(a, b, m, r)x^q \log^w x). \quad (11)$$

Proof of theorem 4. It follows from our conditions and the theory of G – spaces (2) that for all $p \geq 2$

$$|\xi(1)|_p \leq C_1 p^{1/m} \log^{-r/m} p, \quad |\eta|_p \leq C_2 p^{1/a} \log^{-b/a} p.$$

We obtain using the inequality (0) with optimal constant $B(p)$:

$$|S|_p \leq C_3 p^{1+(1/2a)+1/m} \log^{-1-(b/(2a)-(r/m))} p = C_3 p^{1/q} \log^{-w/q} p.$$

The last inequality is equivalent to (11), see (2).

References

- Bagdasarov D.R., Ostrovsky E.I., 1995. Reversion of Chebyshev's Inequality. Probab. Theory Appl., v. 40 **4**, 737 - 742.
- Buldygin V.V., Mushtary D.I., Ostrovsky E.I., Puchalsky M.I., 1992. New Trends in Probability Theory and Statistic. Mokslas; Amsterdam, New York.
- Gine E., Mason D.M., Zaitsev A.Yu., 2003. The L_1 - norm Density Estimation Process. Annals Probab., v. 31, **2**, 719 - 768.
- Gut A., 1988. Stopped Random Walks. Springer Verlag, Berlin - Heidelberg - New York.
- Gut A., 2003. On the Moment Problem for Random Sums. Journal of Appl. Probab., v. 40 *N*^o 3, 707 - 802.
- Hitczenko P., 1990. Best constant in martingale version of Rozental's Inequality. 1990. Annals Probab., v. 18, 1656 - 1668.
- Lesign Emm., Dalibor, 2001. Large deviations for martingales. Stochastic Process. Appl;., **96**, 143 - 159.
- Ostrovsky E.I., 1999. Exponential Estimations for the Random Fields. OINPE, Moscow (in Russian).

Ben - Gurion University, Department of Mathematic.

ISRAEL, Beer - Sheva city, 84105, Ben - Gurion street, 4. P.O.Box
61.

e - mail: Galaostr@cs.bgu.ac.il